Progress of the 1/12° Global HYCOM Effort

E. Joseph Metzger and Alan J. Wallcraft

HYCOM Consortium Meeting 6-8 December 2005 RSMAS, Miami FL

maintaining the data needed, and c including suggestions for reducing	lection of information is estimated to ompleting and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding an DMB control number.	ion of information. Send commentarters Services, Directorate for Inf	ts regarding this burden estimate formation Operations and Reports	or any other aspect of the state of the stat	his collection of information, Highway, Suite 1204, Arlington		
1. REPORT DATE DEC 2005		2. REPORT TYPE			3. DATES COVERED 00-00-2005 to 00-00-2005		
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER			
Progress of the 1/12degree Global HYCOM Effort					5b. GRANT NUMBER		
					5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)					5d. PROJECT NUMBER		
					5e. TASK NUMBER		
					5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory, Stennis Space Center, MS, 39529					8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITO	RING AGENCY NAME(S) A	10. SPONSOR/MONITOR'S ACRONYM(S)					
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)					
12. DISTRIBUTION/AVAIL Approved for publ	LABILITY STATEMENT ic release; distributi	ion unlimited					
13. SUPPLEMENTARY NO	OTES						
14. ABSTRACT							
15. SUBJECT TERMS							
16. SECURITY CLASSIFIC		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF			
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)	20	RESPONSIBLE PERSON		

Report Documentation Page

Form Approved OMB No. 0704-0188

1/12° Global HYCOM: Initial Development and Evaluation

E.J. Metzger, A.J. Wallcraft, H.E. Hurlburt, E.P. Chassignet and W.J. Schmitz, Jr.

Ocean Sciences Meeting 20-24 February 2006 Honolulu, HI


HYCOM Long-term Goals for Operational Ocean Prediction

- 1/12° fully global ocean prediction system transitioned to NAVO in 2007
 - Include shallow water, minimum depth 5 m
 - Coupled sea-ice model (LANL CICE)
- Increase to 1/25° resolution globally by the end of the decade
 - Optimal resolution for basin-scale
 - Boundary conditions for coastal models

Global HYCOM Configuration

- Horizontal grid: 1/12° equatorial resolution
 - 4500 x 3298 grid points, 6.5 km spacing on average, 3.5 km at pole
- Mercator 79°S to 47°N, then Arctic dipole patch
- Vertical coordinate surfaces: 26-28 for σ_0 , 32 for σ_2^*
- KPP and GISS mixed layer models
- Thermodynamic (energy loan) sea-ice model
- Surface forcing: wind stress, wind speed, thermal forcing, precipitation, relaxation to climatological SSS
- Monthly river runoff (986 rivers)
- Initialize from January climatology (GDEM3) T and S, then SSS relaxation from PHC 3.0
 - No subsurface relaxation to climatology

1/12° Global HYCOM snapshot: SSH and ice (gray)



- Running at NAVO under DoD Challenge
- 190K CPU hrs/model year on 784 CPUs


7.2 TB/model year for daily 3-D output

1/12° Global HYCOM Experiments

- ECMWF Reanalysis (ERA15) climatological wind and thermal forcing
 - Annual bias corrections to air temperature (ERA40), radiative fluxes (ISCCP) and precipitation (GPCP)
- σ_0 simulations:
 - 26-layers with KPP for 6 model years
 - 28-layers with GISS for 3 model years
- σ_2^* simulations:
 - 32-layers with GISS for 9 model years


Initial 1/12° Global HYCOM σ₀ Simulation


- Major shortcomings:
 - Poor simulation of both Gulf Stream and Kuroshio
 - Poor representation of tropical current systems
 - Unrealistic transport at key locations:
 - Florida Straits (23 Sv vs. ~32 Sv) [simulated vs. observed]
 - Drake Passage (91 Sv vs. ~134 Sv)
 - Pacific to Indian Ocean Throughflow (22 Sv vs. ~10 Sv)



Improved 1/12° Global HYCOM σ₀ Simulation


- Modifications
 - Added two layers (26 → 28) and changed layer structure
 - Increased eddy viscosity: A = 30 m²/s constant everywhere
 - Increased Smagorinsky diffusion: .05 → .1
 - KPP → GISS






ŏ

Improvements in Tropical Pacific Current Structure




1/12° Global HYCOM Mean Gulf Stream And Kuroshio Pathways



Latest σ_2^* simulation


Mean over four model years

ERA15 climatological wind & thermal forcing




1/12° Global HYCOM σ₂* Simulation

The Atlantic subpolar gyre generally looks good




35W

Global SSH Variability




Oct 92 – Nov 98 SSH variability based on T/P, ERS-1 and ERS-2 altimeters (from Collecte, Localisation, Satellites (CLS))




SSH variability from 1/12° global HYCOM σ₂* with climatological wind and thermal forcing

Mediterranean Sea Outflow Salinity section at 36°N




Atlantic Meridional Overturning Circulation

 σ_0 vs. σ_2^*




Atlantic Meridional Overturning Circulation



Velocity Cross-section Along Luzon Strait


Sb-ADCP data (top) vs. 1/12° global HYCOM (bottom) in the upper 300 m Section along 21°N between 118.5°E and 124.0°E




Sb-ADCP data from Liang et al. (2003, DSR Pt. II)


Mean from HYCOM with ERA15 wind and thermal forcing

No ocean data assimilation in HYCOM






Vertical Structure in the Equatorial Pacific





Deep Flow Through Samoan Passage



Observed mean northward transport below 4000 m = 6.0 Sv Rudnick (1997,JGR)

HYCOM σ_2^* in layers 27-32 = 9.3 Sv





Transport Comparisons at Key Locations

Section	Obs.	Orig. σ_0	Mod. σ_0	Orig. σ_2^*	Mod. σ_2^*
PCM-1	23	24.8	24.7	25.5	26.4
Bering Strait	1	1.1	1.1	1.1	1.1
PACIO TF	-10	-21.8	-24.1	-18.2	-17.0
STACS	30-34	23.2	23.1	22.9	24.0
Yucatan Channel	23-27	23.3	21.6	21.8	22.0
Denmark Strait	-2.9	-2.9	-2.3	-2.3	-2.9
Drake Passage	134	91.3	96.2	152.4	146.4

Modified σ_2^* experiment uses a new topography with sill depth refinements in the IAS, Indonesian Seas, etc.

Future Work (FY06)

- ■Ten May 2001 June 2002 assimilative runs in FY06
 - Time period with three satellite altimeters
 - Five with bi-weekly 30-day forecasts
- Near real-time nowcast/forecast starting in mid-FY06
- Interannual non-assimilative case:
 - 1995-present using NOGAPS
- Coupling with LANL CICE via ESMF

6 Dec 2005 20